



Helping babies breathe since 1998.

Infasurf Dosing Chart

For Intratracheal Administration ONLY

INDICATION

Infasurf is indicated for the prevention of Respiratory Distress Syndrome (RDS) in premature infants at high risk for RDS and for the treatment of premature infants who develop RDS. Infasurf decreases the incidence of RDS, mortality due to RDS, and air leaks associated with RDS.

PROPHYLAXIS

Prophylaxis therapy at birth with Infasurf is indicated for premature infants <29 weeks of gestational age at significant risk for RDS. Infasurf prophylaxis should be administered as soon as possible, preferably within 30 minutes after birth.

TREATMENT

Infasurf therapy is indicated for infants ≤72 hours of age with RDS (confirmed by clinical and radiologic findings) and requiring endotracheal intubation.

Birth Weight (grams)	Total Dose (mL)	Vial(s) Needed*	Birth Weight (grams)	Total Dose (mL)	Vial(s) Needed*
600-650	1.9	3 mL	1301-1350	4.0	6 mL
651-700	2.0	3 mL	1351-1400	4.1	6 mL
701-750	2.2	3 mL	1401-1450	4.3	6 mL
751-800	2.3	3 mL	1451-1500	4.4	6 mL
801-850	2.5	3 mL	1501-1550	4.6	6 mL
851-900	2.6	3 mL	1551-1600	4.7	6 mL
901-950	2.8	3 mL	1601-1650	4.9	6 mL
951-975	2.9	3 mL	1651-1700	5.0	6 mL
976-1000	3.0	3 mL	1701-1750	5.2	6 mL
1001-1050	3.1	6 mL	1751-1800	5.3	6 mL
1051-1100	3.2	6 mL	1801-1850	5.5	6 mL
1101-1150	3.4	6 mL	1851-1900	5.6	6 mL
1151-1200	3.5	6 mL	1901-1950	5.8	6 mL
1201-1250	3.7	6 mL	1951-1975	5.9	6 mL
1251-1300	3.8	6 mL	1976-2000	6.0	6 mL

*Infasurf is available in 3mL and 6mL single-dose vials.

Doses were calculated using the midpoint of each weight range and rounded off to the one decimal point.

CALCULATING DOSE

- Each dose of **Infasurf** is 3 mL/kg body weight at birth
- The **Infasurf Dosing Chart** shows the total dose per administration for a range of birth weights

Please see Indication listed above.
Full Prescribing Information affixed below.

IMPORTANT CONSIDERATIONS

Infasurf should be administered under the supervision of clinicians experienced in the acute care of newborn infants with respiratory failure who require intubation.

ADDITIONAL INFORMATION:

- Store at refrigerated temperature 2° to 8°C (36° to 46°F).
- Infasurf settles during storage—gently swirl or gently invert repeatedly to homogenize settled suspension. **DO NOT SHAKE.**
- Warming not required—if desired, may be warmed to room temperature.
- Never use radiant/artificial warming devices to warm Infasurf.
- Do not remove from refrigerator for more than 24 hours.
- Must not be returned to refrigerator more than once.

Affix PI Here



Manufactured by ONY Biotech
1576 Sweet Home Road | Amherst, NY 14228 | 716.636.9096 | 877.274.4669
infasurf.com



Made in the USA

APM-0073 02/2018